

Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of claims:

1-125. (Canceled)

126. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial cells, the method comprising

(i) contacting the high G+C Gram-positive bacterial cells or dormant, moribund or latent *Mycobacterium tuberculosis* bacterial high G+C Gram-positive bacterial cells *in vitro* with an isolated polypeptide having at least 95% sequence identity with SEQ ID NO:2 or comprising at least 50% sequence identity with amino acid residues 117 to 184 of SEQ ID NO:2, wherein said polypeptide is capable of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating a dormant, moribund, or latent *Mycobacterium tuberculosis* high G+C Gram-positive bacterial cells; and

(ii) incubating the said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial bacterial cells in culture medium containing the polypeptide, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial bacterial cells.

127. (Previously Presented) The method of claim 126, wherein the polypeptide is recombinant.

128. (Currently Amended) The method of claim 126 or 127, wherein said dormant, moribund or latent high G+C Gram-positive bacterial bacterial cells is are present in a sample, and the method identifies a the presence of dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial cells in the sample by detecting growth of high G+C Gram-positive bacterial bacterial cells in the sample.

129-130. (Canceled)

131. (Previously Presented) The method of claim 126 or 127, wherein the polypeptide is in unit dosage form.

132-143. (Canceled)

144. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells, the method comprising

—— (i) contacting the high G+C Gram-positive bacterial cells or dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells *in vitro* with a cell strain expressing a nucleic acid encoding a polypeptide having at least 50% sequence identity with amino acid residues 117 to 184 of SEQ ID NO:2; and

—— SEQ ID NO: 2;

—— a polypeptide having at least 95% sequence identity with SEQ ID NO: 2

—— or a polypeptide comprising at least amino acid residues 117 to 184 of SEQ ID NO: 2;

wherein said polypeptide is capable of resuscitating a dormant, moribund, or latent *Mycobacterium tuberculosis* cell;

—— and

—— (ii) incubating the said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells and the cell strain in culture medium, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial cells.

145-148. (Canceled)

149. (Previously Presented) The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:2.

150. (Previously Presented) The method of claim 126, wherein the isolated polypeptide comprises amino acid residues 117 to 184 of SEQ ID NO:2.

151-156. (Canceled)

157. (Previously Presented) The method of claim 126, wherein the polypeptide is purified essentially to homogeneity.

158. (Cancelled)

159. (Previously Presented) The method of claim 128, wherein the sample is taken from a human or animal.

160. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells, the method comprising

_____ (i) contacting ~~the high G+C Gram-positive bacterial cells or dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells~~ high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells *in vitro* with a purified polypeptide comprising SEQ ID NO:2, wherein said polypeptide is capable of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating a dormant, moribund, or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* cells; and

_____ (ii) incubating ~~the said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial cells~~ in culture medium containing the polypeptide, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial cells.

161. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells, the method comprising

_____ (i) contacting ~~the high G+C Gram-positive bacterial cells or dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells~~ high G+C Gram-positive bacterial *Mycobacterium tuberculosis* bacterial-cells *in vitro* with a

purified polypeptide comprising at least amino acid residues 117 to 184 of SEQ ID NO:-2, wherein said polypeptide is capable of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating a-dormant, moribund, or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis*-cells; and

_____ (ii) incubating ~~the said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells~~ in culture medium containing the polypeptide, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells.

162. (Currently Amended) The method of claim 160 or 161, wherein said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells are present in a sample, and the method identifies a the presence of dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis*-bacterial-cells in the sample by detecting growth of high G+C Gram-positive bacterial bacterial-cells in the sample.

163. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis*-bacterial-cells, the method comprising

_____ (i) contacting ~~the-high G+C Gram-positive bacterial cells or dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells~~ in vitro with a cell strain expressing a nucleic acid encoding a polypeptide comprising SEQ ID NO:-2, wherein said polypeptide is capable of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating a-dormant, moribund, or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis*-cells; and

_____ (ii) incubating ~~the said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells~~ and said cell strain in culture medium, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial bacterial-cells.

164. (Currently Amended) A method of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating dormant, moribund or latent high G+C Gram-positive bacterial *Mycobacterium tuberculosis*-bacterial-cells, the method comprising

_____ (i) contacting ~~the~~ high G+C Gram-positive bacterial cells or dormant, moribund or latent high G+C Gram-positive bacterial ~~bacterial~~-cells *in vitro* with a cell strain expressing a nucleic acid encoding a polypeptide comprising at least amino acid residues 117 to 184 of SEQ ID NO: 2, wherein said polypeptide is capable of stimulating growth of high G+C Gram-positive bacterial cells or of resuscitating a ~~dormant, moribund, or latent high G+C Gram-positive bacterial~~ *Mycobacterium tuberculosis*-cells, and

_____ (ii) incubating ~~the~~ said high G+C Gram-positive bacterial cells or said dormant, moribund or latent high G+C Gram-positive bacterial ~~bacterial~~-cells and ~~said~~ cell strain in culture medium, thereby stimulating growth of said high G+C Gram-positive bacterial cells or resuscitating said dormant, moribund or latent high G+C Gram-positive bacterial ~~bacterial~~-cells.

165. (New) The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:36 or SEQ ID NO:43.

166. (New) The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:7.

167. (New) The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:2.

168. (New) The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:3.

169. The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:4.

170. The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:5.

171. The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:6.

172. The method of claim 126, wherein the isolated polypeptide comprises SEQ ID NO:8.